

Opportunistic Feeding of an Adult Humpback Whale (*Megaptera novaeangliae*) Migrating Along the Coast of Southeastern Queensland, Australia

Karen A. Stockin¹ and Elizabeth A. Burgess²

¹Coastal-Marine Research Group, Institute of Natural Resources, Massey University,
Private Bag 102 904, North Shore MSC, Auckland, New Zealand

²Tangalooma Marine Education and Research Centre, P.O. Box 1200, Eagle Farm Queensland 4009, Australia

Abstract

In the austral winter of July 2004, off southeastern Queensland, Australia, we observed apparent feeding by an adult humpback whale (*Megaptera novaeangliae*) in association with Indo-Pacific bottlenose dolphins (*Tursiops aduncus*). This is the first documented case of an adult humpback whale from the southern hemisphere Group V stock feeding along the migratory corridor. This observation also represents the first published record of a humpback whale feeding in association with Indo-Pacific bottlenose dolphins.

Key Words: humpback whale, *Megaptera novaeangliae*, Indo-Pacific bottlenose dolphin, *Tursiops aduncus*, opportunistic feeding, migration, Australia

Introduction

Like most species of mysticetes, humpback whales (*Megaptera novaeangliae*) undertake annual migrations from high latitude feeding areas to low latitude breeding areas. Between June and November each year, humpback whales travel along the east coast of Australia (Paterson & Paterson, 1989) during their annual migration between the feeding grounds off Antarctica (Bryden, 1978; Chittleborough, 1959) and the breeding grounds off the Great Barrier Reef (Paterson & Paterson, 1989; Simmons & Marsh, 1986). Waters adjacent to Cape Moreton (27° 02' S, 153° 28' E) off southeastern Queensland, are part of the migration route for so-called Area V (130° E-170° W) humpback whales during the austral winter months (Paterson, 1991). Herein, we report an opportunistic observation of apparent feeding involving an adult humpback whale and a group of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) during this migration.

Materials and Methods

Observations were made from the *M.V. Tangalooma Jet*, a 40 m jet-propelled commercial whale-watching catamaran, which conducts tours from Moreton Island, focusing primarily on humpback whales during their migration. Data presented are the result of an observation made by the authors during opportunistic whale watching. Photographs were taken using a Nikon D100 digital SLR and a Nikon ED AF 70-300 mm lens.

Results

On 5 July 2004, observations were made of apparent feeding by a humpback whale and a group of adult Indo-Pacific bottlenose dolphins. At 1250 h, three humpbacks were encountered within 200 m of each other during their northern migration towards the breeding grounds. They were approximately 1.5 km off Cape Moreton, the northeastern headland of Moreton Island. At 1310 h, a group of approximately 50 to 60 Indo-Pacific bottlenose dolphins were observed heading east towards the whale watch vessel. They were identified as Indo-Pacific bottlenose dolphins based on a long rostrum and overall morphology. The dolphins initially appeared as a single group, travelling "line abreast" (Neumann & Orams, 2003, p. 143), on course towards the travelling whales. Within approximately 200 m of the vessel, the dolphins divided into two groups, with one subgroup of approximately 30 to 40 animals turning away to the northwest of the whales and the vessel. The remaining subgroup of approximately 15 to 20 adult dolphins passed within 5 m of the bow of the vessel. At 1316 h, this subgroup began to engage in active foraging behaviour; high-speed pursuits by individual dolphins (as defined in Neumann & Orams, 2003) were noted on several occasions, and successful prey capture by dolphins was observed in two instances. "Carouseling" (Neumann & Orams, 2003) and "kerplunking" (Connor et al.,

2000; Nowacek, 1999) also were observed. The prey was not identified to species, but clearly was a type of small schooling baitfish, approximately 12 cm in length.

At 1327 h, one of the humpback whales adjusted course to the northeast and separated from the other two whales, which remained within 300 m of each other as they continued to travel in a north northeast direction. The solitary whale of unknown gender was estimated to be approximately 14 m in length compared to the whale watch vessel. This whale then approached the subgroup of dolphins. The dolphins were foraging and were continuously observed in close association with a prey "baitball," which was clearly visible at the water surface. At 1330 h, in water of 23 m depth, and at a location 27° 02' S and 153° 28' E, the whale entered the immediate vicinity of the dolphins (Figure 1) and was observed rolling laterally through the fish baitball in the centre of foraging dolphins.

Evidence of the whale feeding included clear expansion of the ventral pleats (Figure 2) and subsequent contraction, its partially open mouth closing just under the surface of the water, and baitfish breaking the surface near the whale. The whale continued travelling in close proximity (i.e., within two whale body lengths; see Figure 1) to the dolphins for a further 11 min after the described observation. Although no further baitballs were observed, it cannot be dismissed that the whale continued to feed on loose aggregations of the baitfish because the dolphins continued to forage during this period. From this time forward, the distance between the whale and the dolphins could not be closely observed because the whale watch vessel began to depart the area; subsequently, the encounter terminated at 1342 h. No photo-identification was possible because the whale did not fluke up at any point during the encounter.

Discussion

Although sightings of humpback whales in close proximity to bottlenose dolphins have been previously noted (e.g., Weller et al., 1996), they are not common and no previously published literature relates to interactions with Indo-Pacific bottlenose dolphins. Furthermore, feeding interactions between these two species have not been reported previously. It appeared as if the humpback whale detected the bottlenose dolphins actively foraging and subsequently fed from the same baitball. The Indo-Pacific bottlenose dolphins appeared to work cooperatively to concentrate a school of baitfish, which were clearly visible both at and just below



Figure 1. Humpback whale observed in close association with foraging Indo-Pacific bottlenose dolphin off Cape Moreton, southeastern Queensland, in July 2004 (Photograph from K. A. Stockin)



Figure 2. View of expanding ventral throat pleats of a humpback whale feeding on baitfish off Cape Moreton, southeastern Queensland, in July 2004 (Photograph from K. A. Stockin)

the surface of the water throughout the observation period.

Few records of humpback whales feeding during their migration, or on the breeding grounds, exist (Baraff et al., 1991; Gendron & Urbán, 1993; Salden, 1989; Stone et al., 1987; Swingle et al., 1993). To date, observations relate only to the northern hemisphere, and have involved juvenile animals. In the low latitude waters of the southern hemisphere, it is believed that humpback whales do not generally feed during migration (Chittleborough, 1965; Dawbin, 1966). Dall & Dunstan (1957) recorded food present in only one of over 2,000 humpback whales examined at Tangalooma whaling station. Chittleborough (1965) suggested that humpback whales seldom find sufficiently dense swarms of plankton off the Australian coast to stimulate feeding but, instead, subsist on stored fat reserves. This subsistence period is clearly evident in the progressive decline in oil yields obtained from low latitude whaling stations.

This is the first documented evidence of opportunistic feeding by an adult humpback whale during the southern breeding migration. Despite the absence of vertical or subsurface

lunge-feeding (Baraff et al., 1991; Salden, 1989), bubble clouds (Hain et al., 1982; Jurasz & Jurasz, 1979; Weinrich et al., 1992), or defecation (Gendron & Urbán, 1993), apparently the whale was foraging, based on extended then contracted throat pleats, open mouth, and the presence of baitfish.

The diet of humpback whales in the northern hemisphere (Hain et al., 1982) typically consists of small schooling fish such as herring (*Clupea harengus*), capelin (*Mallotus villisus*), and sand lance (*Ammodytes* sp.), although humpback whales in the waters off southeastern Alaska are known to exhibit a more diverse diet (Sharpe, 2001). In the southern hemisphere, humpback whales feed predominantly on euphausiids (e.g., *Euphausia superba*, *E. spinifera*, and copepods, *Calanus propinquus*) (Winn & Reichley, 1985). During the observation reported here, it was apparent that the whale was feeding directly on baitfish and while the species could not be confirmed, our observations suggest they were most likely sardines (*Sardinops sagax*). This is consistent with reports of various species of pelagic baitfish shoaling close to the southern Queensland coastline, including sardine, round eye herring (*Etrumeus teres*), blue mackerel (*Scomber australasicus*), and yellowtail scad (*Trachurus novaezelandiae*) (J. Staunton-Smith, pers. comm.).

We make no claim that humpback whales regularly undertake opportunistic feeding during their journey along migratory corridors, or on breeding grounds. It is not possible to ascertain how frequently such opportunistic foraging occurs; however, if common, we would expect this behaviour to have been previously reported, given the observation efforts through dedicated research and frequent whale watching worldwide. Baraff et al. (1991) suggested that occasional opportunistic feeding may be a mechanism undertaken to offset any energy deficiency incurred through the winter. Length at sexual maturity for humpback whales in Australian waters is estimated to be 11.2 m and 11.7 m for males and females, respectively (Chittleborough, 1955a; Chittleborough, 1955b). The length of the whale in the present observation implies it was an adult and not a juvenile or subadult as documented in previous northern hemisphere examples (Baraff et al., 1991; Salden, 1989; Swingle et al., 1993). The significance of this remains unclear, although it suggests that this type of opportunistic behaviour may not be exclusive to immature animals.

Acknowledgments

The authors thank the staff at Tangalooma Wild Dolphin Resort, Queensland, especially Trevor Hassard, Ben Carroll, and the skipper Jeff Osborne

and crew of the *M.V. Tangalooma Jet*. Special thanks are extended to Phil Clapham and Mike Noad for their comments on our observations and to Phil Clapham, Scott Baker, Dan Salden, Mark Orams, and Ingrid Visser for their comments on earlier drafts of this manuscript. KAS is supported by a Commonwealth Scholarship.

Literature Cited

- Baraff, L. S., Clapham, P. J., Mattila, D. K., & Bowman, R. S. (1991). Feeding behavior of a humpback whale in low-latitude waters. *Marine Mammal Science*, 7, 197-202.
- Bryden, M. M. (1978). Whales and whaling in Queensland waters. *Proceedings of the Royal Society of Queensland*, 88, 5-18.
- Chittleborough, R. G. (1955a). Aspects of reproduction in the male humpback whale, *Megaptera nodosa* (Bonnaterre). *Australian Journal of Marine and Freshwater Research*, 6, 1-29.
- Chittleborough, R. G. (1955b). Puberty, physical maturity, and relative growth of the female humpback whale, *Megaptera nodosa* (Bonnaterre), on the western Australian coast. *Australian Journal of Marine and Freshwater Research*, 6, 315-327.
- Chittleborough, R. G. (1959). Australian marking of humpback whales. *Norsk Hvalfangst-Tidende*, 48, 47-55.
- Chittleborough, R. G. (1965). Dynamics of two populations of the humpback whale, *Megaptera novaeangliae* (Borowski). *Australian Journal of Marine and Freshwater Research*, 16, 33-128.
- Connor, R. C., Heithaus, M. R., Berggren, P., & Miskis, J. L. (2000). "Kerplunking": Surface fluke splashes during shallow-water bottom foraging by bottlenose dolphins. *Marine Mammal Science*, 16, 646-653.
- Dall, W., & Dunstan, D. J. (1957). Krill (*Euphausia superba* Dana) from a humpback whale, *Megaptera nodosa* (Bonnaterre), caught off southern Queensland. *Norsk Hvalfangst-Tidende*, 46, 6-9.
- Dawbin, D. H. (1966). The seasonal migratory cycle of humpback whales. In K. S. Norris (Ed.), *Whale, dolphins and porpoises* (pp. 145-170). Berkeley: University of California Press.
- Gendron, D., & Urbán, R. (1993). Evidence of feeding by humpback whales (*Megaptera novaeangliae*) in the Baja California breeding ground, Mexico. *Marine Mammal Science*, 9, 76-81.
- Hain, J. H. W., Carter, G. S., Kraus, S. D., Mayo, C. A., & Winn, H. E. (1982). Feeding behavior of the humpback whale, *Megaptera novaeangliae*, in the western North Atlantic. *Fishery Bulletin*, 80, 259-268.
- Jurasz, C. M., & Jurasz, V. P. (1979). Feeding modes of the humpback whale, *Megaptera novaeangliae*, in south-east Alaska. *Scientific Reports of the Whales Research Institute*, 31, 69-83.

- Neumann, D. R., & Orams, M. B. (2003). Feeding behaviours of short-beaked common dolphins, *Delphinus delphis*, in New Zealand. *Aquatic Mammals*, 29, 137-149.
- Nowacek, D. (1999). *Sound use, sequential behavior and the ecology of foraging bottlenose dolphins* (*Tursiops truncatus*). Doctoral dissertation, Woods Hole Oceanographic Institute, Woods Hole, MA.
- Paterson, R. A. (1991). The migration of humpback whales *Megaptera novaeangliae* in east Australian waters. *Memoirs of the Queensland Museum*, 29, 333-341.
- Paterson, R., & Paterson, P. (1989). The status of the recovering stock of humpback whales *Megaptera novaeangliae* in east Australian waters. *Biological Conservation*, 47, 33-48.
- Salden, D. R. (1989). An observation of apparent feeding by a sub-adult humpback whale off Maui, Hawaii. *Abstracts of the Eighth Biennial Conference on the Biology of Marine Mammals* (p. 58), Pacific Grove, California.
- Sharpe, F. A. (2001). *Social foraging of the southeast Alaskan humpback whale*. Doctoral dissertation, Simon Fraser University, Burnaby, BC, Canada.
- Simmons, M. L., & Marsh, H. (1986). Sightings of humpback whales in Great Barrier Reef waters. *Scientific Reports of the Whales Research Institute*, 37, 31-46.
- Stone, G. S., Katona, S. K., & Tucker, E. B. (1987). History, migration and present status of humpback whales *Megaptera novaeangliae* at Bermuda. *Biological Conservation*, 42, 133-145.
- Swingle, W. M., Barco, S. G., Pitchford, T. D., McLellan, W. A., & Pabst, D. A. (1993). Appearance of juvenile humpback whales feeding in the nearshore waters of Virginia. *Marine Mammal Science*, 9, 309-315.
- Weinrich, M. T., Schilling, M. R., & Belt, C. R. (1992). Evidence for acquisition of a novel feeding behaviour: Lobtail feeding in humpback whales, *Megaptera novaeangliae*. *Animal Behaviour*, 44, 1059-1072.
- Weller, D. W., Schiro, A. J., Cockcroft, V. G., & Wang, D. (1996). First account of a humpback whale (*Megaptera novaeangliae*) in Texas waters, with a re-evaluation of historical records from the Gulf of Mexico. *Marine Mammal Science*, 12, 133-137.
- Winn, H. E., & Reichley, N. E. (1985). Humpback whale *Megaptera novaeangliae* (Borowski, 1781). In S. H. Ridgway & R. J. Harrison (Eds.), *Handbook of marine mammals, Volume 3* (pp. 241-273). London: Academic Press.